Supplementary Table 1. Distribution of patients according to underlying etiology

	Patient (n)	Patient %
Postinfectious cough	59	37.8
Asthma	39	25
COPD	11	7.1
Gastroesophageal reflux disease	11	5.8
Hypersensitivity pneumonitis	7	4.5
Allergic cough	5	3.2
Unclassifiable ILD	5	3.2
ACE inhibitor use	4	2.6
CTD-ILD	3	1.9
Asthma–COPD overlap syndrome	2	1.3
Bronchiectasis	2	1.3
Nasal polyposis	2	1.3
Pre-COPD	2	1.3
Idiopathic pulmonary fibrosis	1	0.6
Obesity hypoventilation syndrome	1	0.6
Pleuroparenchymal fibroelastosis	1	0.6
Sarcoidosis	1	0.6

COPD: Chronic obstructive pulmonary disease, CTD-ILD: Connective tissue disease associated interstitial lung disease, ILD: Interstitial lung disease, ACE: Angiotensin-converting enzyme

Supplementary Table 2. Mean spirometric z-scores of major diagnostic subgroups in patients with subacute cough

	Postinfectious cough (n = 59)	Obstructive lung diseases (n = 56)	ILDs and others (n = 40)
FEV_1	-0.85±1.01	-1.42±1.11	-0.87±1.08
FVC	-1.22±1.21	-1.48±1.21	-1.20±1.02
FEV_1/FVC	0.98±1.11	0.21±1.50	0.68±1.43
$FEF_{25–75}$	0.12±0.84	-1.67±1.12	0.08±1.14

FEF_{25-75}: Forced expiratory flow between 25% and 75% of FVC, FEV_1: Forced expiratory volume in one second, FVC: Forced vital capacity, ILDs: Interstitial lung diseases